

DC's Real-Time HIV Data Reveal Segregation, Options for Improving Care

Kenya has been using similar technology for three years. DC's efforts are among the first in the United States.

January 8, 2021 By [Heather Boerner](#)

A new study offers ways to help Washington, DC, residents realize the health benefits of [Undetectable = Untransmittable](#). Treatment adherence support and other HIV programs should focus on hard-hit groups in hot-spot neighborhoods, and public health officials should invite community participation in designing locally relevant programs.

In particular, health officials should focus on getting treatment adherence messages to Black and Latino same-gender-loving men in central DC, work with schools to decrease truancy and increase engagement in care in south and southeastern DC, and target existing programs that help people under age 26 access health insurance in central and southeast DC.

Washington, DC, ward map Courtesy of League of Women Voters

These are the findings of a DC Department of Health modeling study, based on 2018 data. The analysis found that in the years leading up to 2018, rates of viral suppression worsened among residents of wards seven and eight who didn't acquire HIV through gay sex. Among men who have sex with men, viral suppression plateaued in the years leading up to and including 2018.

"DC so far has implemented homogeneous policies on young people across the city," Suparna Das, PhD, a health department statistician, and colleagues wrote in the journal [Clinical Infectious Diseases](#).

But the analysis indicates that those efforts, including social media campaigns and attempts to

help young people access health insurance separate from their parents, should instead focus on hot spots. “This analysis should be used for data-to-care planning activities,” the authors recommended.

Courtesy of AIDSvu.org

HIV mapping projects aren’t new. [AIDSvu.org](#) has been providing granular data on viral suppression [for years](#). And in 2019, amfAR, The Foundation for AIDS Research, launched the [Ending the HIV Epidemic database](#). Kenya has used real-time data to target HIV prevention resources since 2016, according to a [Medscape report](#).

Traditional methods of finding communities with low levels of viral suppression rely on person-to-person contact tracing, which can be slow, inefficient and incomplete.

But the launch of [Ending the HIV Epidemic: A Plan for America](#) in 2019 came with the directive to use snippets of HIV DNA from blood tests, known as molecular tracing, to track HIV outbreaks and target resources to hot spots where people are less likely to be living well with HIV. Some activists working toward ending HIV nondisclosure and other criminalization laws have demanded that this kind of real-time tracking be removed from the plan, citing the potential that the data could be used to [prosecute people living with HIV](#).

This model may provide an alternative to molecular tracing. It culls information on HIV diagnoses and viral loads from the enhanced HIV/AIDS Reporting System.

The model showed that while people with HIV live throughout DC, same-gender-loving men without viral suppression are more likely to live in the city’s central wards one, two and five. People who didn’t acquire HIV through gay sex, meanwhile, are more likely to have a detectable viral load and to be disengaged from care in wards seven and eight. They are also more likely to

be Black, more likely to live in poverty and less likely to have access to the internet. These findings are consistent with the stark racial segregation in the nation's capital.

The findings particularly call out young people ages 12 to 18 outside of gay male hot spots, gay men ages 21 to 24 within those gay hot spots and people who inject drugs as being at highest risk for having a detectable viral load.

These data, the study authors wrote, suggest that existing programs aimed at improving viral suppression should focus messaging and resources within these hot spots. They also suggest that public health officials consider new programs to address the models' findings. These include:

- Targeting treatment adherence activities for same-gender-loving Black and Latino men to gay male hot spots in wards one, two and five;
- Implementing programs within DC's public and charter schools to support adherence;
- Working with parents and schools, colleges and universities in these hot spots to ameliorate truancy and improve HIV treatment adherence;
- Evaluating the digital divide in DC, with an eye to whether internet and mobile phone adherence efforts will be as effective in the city's seventh and eighth wards as it may be in wards one, two and five.

"Overwhelming, clinical evidence has recognized that for HIV, Undetectable = Untransmittable," wrote Das and colleagues. "Thus, it becomes imperative for health departments in collaboration with the community help sustain the undetectable viral load of [people living with HIV]."

[Click here](#) to read the full analysis.