

Delivering Oxygen to Tumors May Be Key in Overcoming Radiation Resistance

Using a targeted approach, researchers slowed head and neck cancer growth in mice.

July 28, 2023 By National Institutes of Health

For most of our tissues and cells, a lack of oxygen, or hypoxia, is bad news. Cancer cells, on the other hand, can thrive in these hypoxic conditions, which render tumors less susceptible to anti-cancer treatments, including radiation. Now, new research may offer a way to break through cancer's hypoxia-induced defenses.

A team led by researchers at Thomas Jefferson University and Drexel University has devised a strategy that combines ultrasound with microbubbles to deliver oxygen and cancer drugs to tumors. The results of [a study published in the journal *Pharmaceutics*](#) indicate that the method — tested in mice — primed tumors to be more vulnerable to radiation therapy, resulting in slowed tumor growth and increased survival. The study authors suggest that, with further development, their approach could be used to enhance the effectiveness of radiation therapy in the clinic for a variety of cancers.

"At the moment there's very little we can do on the clinical side to overcome hypoxia for radiation therapy, despite it being a well-known and documented limitation," said study author John Eisenbrey, PhD, an associate professor of radiology at Thomas Jefferson University. "The hope is that a more localized, more aggressive targeting of hypoxia, like what we've done here, will overcome some of that."

A long-standing staple of cancer management, radiation therapy, also called radiotherapy, is used to treat [more than half](#) of all patients with cancer. With this method, clinicians use high doses of radiation to kill cancer cells.

But many tumors are resistant to radiotherapy, and hypoxia is one of the major reasons why, Eisenbrey said.

"Whenever I say we're delivering oxygen to tumors, a lot of people say, 'Well that's crazy. You're going to encourage them to grow,'" said study co-author Margaret Wheatley, PhD, a professor of biomedical engineering at Drexel University. But radiation therapy, which prevents cancer cells

from replicating, requires an ample supply of oxygen close by to be effective, she explained.

Seeking to beat radiotherapy resistance, Wheatley, Eisenbrey, and their colleagues designed a method that sensitizes tumors to radiotherapy using tiny, delicate bubbles (called microbubbles) that are about the size of red blood cells.

To increase oxygen levels in tumors, they packaged oxygen molecules into the microbubbles. And to prevent cancer cells from consuming the oxygen for their own benefit, they also added the drug lonidamine, which interrupts cancer cell metabolism.

After assembly, the microbubbles are injected and circulate throughout the body, only releasing their payload at the intended target, sparing healthy tissues. The trick to the targeted release is ultrasound. By specifically casting ultrasound waves at a tumor, the researchers can cause the microbubbles that have traveled there to pop and release their contents.

In [a previous study](#), the researchers used the method to treat mice with tumors derived from breast cancer cells, producing favorable results. They upped the ante in the new work, instead working against head and neck cancer cell-derived tumors.

“They are notoriously more hypoxic than other solid tumors and also relatively resistant to radiation therapy,” Eisenbrey said.

The study authors injected mice with microbubbles containing oxygen and lonidamine, bursting them at the tumor site with ultrasound, while also administering a drug called metformin — known to slow cancer growth — orally. Other groups of mice were given less than the full treatment, not receiving one or many of the elements of the strategy prior to radiotherapy.

After exposing the animals to radiation, the researchers tracked the size of the tumors over the following three months.

All groups that received some degree of treatment saw varying levels of slowed tumor growth. Mice treated with fully loaded bubbles and metformin prior to radiotherapy lived the longest, their tumor growth slowing drastically. The authors suggest the reason why is that the increase in oxygen within the tumors made them more vulnerable to radiation treatment.

Gaining ground on head and neck cancer is a very encouraging result, even in mice, Eisenbrey said. However, the team intends to continue fine-tuning the method by optimizing radiation and drug doses for better results and streamlining microbubble fabrication, ultimately bringing their solution closer to reality for patients with cancer.

“The authors are using ultrasound and microbubbles in an innovative way, delivering oxygen and drugs to tackle a very serious clinical dilemma. They obtained some promising results, so I’m curious to see how this approach fares with additional development,” said Randy King, PhD, director of the ultrasound program in the Division of Applied Science and Technology at the National Institute of Biomedical Imaging and Bioengineering (NIBIB).

This research was funded by a grant from NIBIB (R01EB026881).

This Science Highlight describes a basic research finding. Basic research increases our understanding of human behavior and biology, which is foundational to advancing new and better ways to prevent, diagnose, and treat disease. Science is an unpredictable and incremental process—each research advance builds on past discoveries, often in unexpected ways. Most clinical advances would not be possible without the knowledge of fundamental basic research.

This [news release](#) was published by theNational Institute of Biomedical Imaging and Bioengineering on July 10, 2023.

© 2026 Smart + Strong All Rights Reserved.

<http://beta.docker.tusaludmag.com/article/delivering-oxygen-tumors-may-key-overcoming-radiation-therapy-resistance>