TuSalud

Even Lean People With HIV Can Have Fatty Liver Disease

Relying on BMI alone may fail to identify people with liver fat accumulation.

August 11, 2026 By Liz Highleyman

Lean people who are classified as normal weight according to body mass index (BMI) may still develop fatty liver disease and its complications—and those living with HIV may be at higher risk—according to study findings presented at the 26th International AIDS Conference (AIDS 2026) in Rio de Janeiro.

“Normal BMI does not exclude clinically relevant liver disease in HIV,” said presenter Florence Bascombe, RN, a senior research nurse at University College London. “Reliance on body size alone may miss people at risk, highlighting the need for better phenotyping and more nuanced metabolic assessment in HIV care.”

Steatotic liver disease (SLD) is the recently adopted name for fat accumulation in the liver. It includes metabolic dysfunction-associated steatotic liver disease (MASLD), alcohol-related liver disease (ALD) and liver disease attributable to both metabolic factors and alcohol (MetALD).

The new terminology recognizes that SLD is a metabolic condition. Over time, the buildup of fat in the liver (hepatic steatosis) can lead to fibrosis, cirrhosis and liver cancer. Some research finds that lean people with SLD are at greater risk for these complications, as well as higher mortality, despite fewer metabolic abnormalities. Now that hepatitis B can be prevented with a vaccine and hepatitis C can be cured with direct-acting antiviral therapy, SLD is responsible for a growing share of advanced liver disease worldwide.

Fatty liver disease is often associated with obesity, but not always. Studies of the general population find that around 10% of people with normal or low body weight may have SLD, and some research suggests this may rise to 25% among people living with HIV. Focusing on BMI alone can underestimate the prevalence of fatty liver disease in lean people with or without HIV.

To learn more about fatty liver disease in people with HIV, Bascombe and colleagues compared clinical characteristics of HIV-positive people with SLD by BMI category. This cross-sectional analysis used clinical data collected at a large HIV service in central London between 2019 and 2025.

SLD was determined using transient elastography, or FibroScan, a noninvasive ultrasound imaging method that estimates both steatosis (by controlled attenuation parameter) and fibrosis (by liver stiffness measurement). Participants with a BMI below 25 were classified as having lean SLD. A BMI of 25 is the traditional threshold for overweight, while 30 is the cut-off for obesity. The researchers also collected data on liver disease biomarkers, HIV-related factors and cardiometabolic health.

Among 391 people with HIV and fatty liver disease, 57 (15%) were classified as having lean SLD. Compared to people whose BMI was 25 or higher, lean people with SLD were older (median 58 versus 54 years) and generally had more a favorable cardiometabolic profile. Nonetheless, lean and non-lean people with SLD had similar markers of liver disease severity, including liver fibrosis scores.

“Although lean participants appeared metabolically healthier and had slightly less hepatic fat accumulation, we found no evidence that they had a lower burden of clinically relevant liver disease,” Boscombe said.

Of note, hepatitis B coinfection was more common among lean people with SLD in a single-factor analysis. In fact, people with lean SLD were more than twice as likely to have hepatitis B than those with overweight or obesity (19% versus 8%).

Lean SLD was most strongly associated with a history of lipodystrophy, or abnormal body fat distribution. While 11% of people with lean SLD had ever had lipodystrophy, this fell to just 2% among those with overweight or obesity.

Once a hallmark of HIV, people with lipodystrophy may have excess abdominal fat along with fat loss in the face and limbs (lipoatrophy). Although its causes are not fully understood, some early antiretroviral drugs—in particular thymidine analogues such as AZT and early protease inhibitors—were linked to either lipoatrophy or mixed fat gain and loss. In this analysis, lean SLD was associated with use of tenofovir alafenamide (a component of several combination pills including Descovy and Biktarvy).

“Collectively, these findings raise the possibility that lean SLD may not simply represent obesity-driven disease occurring at a lower BMI, but instead may arise through different pathways leading to SLD in people with HIV, including altered fat distribution, adipose dysfunction or potentially mitochondrial mechanisms,” the researchers suggested.

These findings show that screening strategies based on BMI and metabolic factors may miss a subgroup of people at risk for fatty liver disease. This has implications for clinical management, including prescribing of GLP-1 weight-loss medications. One of these, Wegovy (semaglutide) was approved last year for the treatment of metabolic dysfunction-associated steatohepatitis (MASH), the advanced stage of MASLD.

Recognizing this discrepancy, the field is moving away from reliance on BMI alone and toward including body composition measures such as waist circumference, waist-to-height ratio, body

roundness index and body fat percentage.

Click here for more news about HIV and liver disease.

Click here for more reports from AIDS 2026.
POZ is an official media partner of AIDS 2026.

© 2026 Smart + Strong All Rights Reserved.
http://beta.docker.tusaludmag.com/article/even-lean-people-hiv-can-fatty-liver-disease