

Lynparza Shows Long-Term Survival Benefits for Patients With High-Risk BRCA-Positive Breast Cancer

Women with high-risk HER2-negative breast cancer who received PARP inhibitor after standard therapy had improved survival at six years.

December 11, 2024 By Dana-Farber Cancer Institute

Patients with high-risk, BRCA-positive breast cancer who received olaparib [Lynparza] after standard treatment continued to have better survival outcomes than those who received a placebo after a median follow-up of 6.1 years, according to the latest results from the Phase III OlympiA clinical trial presented at the San Antonio Breast Cancer Symposium (SABCS 2024). Results were also published in [The New England Journal of Medicine.](#)]

“The OlympiA trial examines adding one year of the oral PARP inhibitor olaparib after completion of standard treatment for higher risk breast cancer in individuals with pathogenic germline BRCA variants,” said [Judy E. Garber, MD, MPH](#), chief of the Division of Cancer Genetics and Prevention at Dana-Farber Cancer Institute and the presenter of the study.

OlympiA is a multicenter, double-blind study in which 1,836 patients with BRCA-positive, HER2-negative breast cancer were randomly assigned (1:1) to receive either olaparib or a placebo for one year following completion of chemotherapy, surgery, and radiation. Based on previous results from the trial, Olaparib was approved in 2022 in the adjuvant setting for certain patients with HER2-negative, BRCA-positive breast cancer.

“Data from the third prespecified interim analysis presented here provides further support for olaparib’s benefits in patients with high-risk, HER2-negative breast cancer with germline mutations in BRCA1 or BRCA2,” said Garber.

After a median follow-up of 6.1 years, patients treated with olaparib continued to show significant improvements in survival outcomes, in both triple-negative breast cancer as well as estrogen receptor (ER)-positive breast cancer subgroups. Both of the study’s primary endpoints—invasive disease-free survival and distant disease-free survival—were achieved, with olaparib reducing the risk of each by 35%.

At six years, 79.6% of olaparib-treated patients remained free of invasive recurrence and 83.5%

remained free of distant recurrence, compared to 70.3% and 75.7%, respectively, in the placebo group.

Additionally, adjuvant PARP inhibition was associated with a 28% reduction in risk of death, and no increase in the risk of developing myelodysplastic syndrome (MDS) or acute myeloid leukemia (AML), both of which can result from rare complications associated with breast cancer treatment.

“The ongoing data from the OlympiA trial are reassuring in the observations of persistent and increasing benefits for these breast cancer patients in the follow-up phases, improving not only protection against recurrence but also overall survival as well,” Garber said. “This demonstration of efficacy makes it more important than ever that we be able to identify individuals who might benefit when they begin their treatment so that we can plan to introduce olaparib to their care at the most opportune moments.”

There were fewer secondary BRCA-associated cancers reported in the olaparib group—38 versus 57 in the placebo group—but Garber noted that further data and analyses are needed to draw firm conclusions.

There were equal numbers of pregnancies in both groups, which is encouraging and highlights the younger age of this cohort, she added.

“These data highlight the safety of olaparib and, therefore, the possibility of moving PARP inhibitors to the treatment of BRCA-associated breast cancers that are lower risk,” noted Garber. “It also allows us to consider the very preliminary possibility of a safe and effective oral agent that could be developed for cancer interception—to be given intermittently to eliminate cells in BRCA mutation carriers that have begun to transform toward several types of malignancy.”

One limitation of the study was that not all subsets were equally large, but efficacy continues to be demonstrated in all subsets, which is very reassuring, Garber noted.

This study was funded by AstraZeneca and Merck and Co., Inc.

This [news release](#) was published by Dana-Farber Cancer Institute on December 11, 2024.

Click here for more news from [SABCS 2024](#).